

No part of the candidate evidence in this exemplar material may be presented in an external assessment for the purpose of gaining credits towards an NCEA qualification.

2

91262



912620



NEW ZEALAND QUALIFICATIONS AUTHORITY
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

SUPERVISOR'S USE ONLY

Level 2 Mathematics and Statistics, 2016

91262 Apply calculus methods in solving problems

9.30 a.m. Thursday 24 November 2016
Credits: Five

Achievement	Achievement with Merit	Achievement with Excellence
Apply calculus methods in solving problems.	Apply calculus methods, using relational thinking, in solving problems.	Apply calculus methods, using extended abstract thinking, in solving problems.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should attempt ALL the questions in this booklet.

Make sure that you have Formulae Sheet L2–MATHF.

Show ALL working.

If you need more space for any answer, use the page(s) provided at the back of this booklet and clearly number the question.

You must show the use of calculus in answering all questions in this paper.

Check that this booklet has pages 2–12 in the correct order and that none of these pages is blank.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.

Achievement

TOTAL

10

ASSESSOR'S USE ONLY

QUESTION ONE

- (a) A function f is given by $f(x) = 4x^3 - 7x^2 + 2x - 4$.

Find the gradient of the graph of the function at the point where $x = 2$.

$$f(x) = 4x^3 - 7x^2 + 2x - 4 \rightarrow f'(x) = 22$$

$$f'(x) = 12x^2 - 14x + 2$$

$$= 12(2)^2 - 14(2) + 2$$

$$\text{Gradient} = 22$$

- (b) The line $y = x + 3.25$ is a tangent to the graph of the function $f(x) = 3x^2 - 2x + 4$.

Use calculus to show that the line is a tangent to the curve, and that the point where this tangent touches the curve is $(0.5, 3.75)$.

$$y = mx + c$$

$$f'(x) = 6x - 2$$

$$1 = 6x - 2$$

$$= 6(0.5) - 2$$

$$3 = 6x$$

$$m = 1$$

$$0.5 = x$$

- (c) The function $f(x) = 2x^3 + kx^2 + 5$ has a minimum turning point when $x = 1$.

What are the coordinates of the **maximum** turning point?

$$f(x) = 2x^3 + kx^2 + 5$$

$$f'(x) = 6x^2 + k$$

$$= 6(1)^2 + k$$

$$-6 = m$$

$$-6 = (6(1)^2) + k$$

$$0 = 1 + k$$

$$-1 = k$$

- (d) The equation of a function $y = f(x)$ has gradient function of the form $f'(x) = 2x - a$, where a is a constant.

The point $(3, 4)$ is the turning point on the graph of the function.

Find the equation of the function.

$$f'(x) = 2(3) - a$$

$$= 6 - a$$

$$f(x) = 6x - ax$$

$$f(x) = 6x - 4.67a$$

$$4 = 6x - ax$$

$$4 = 6(3) - a(3)$$

$$4 = 18 - a(3)$$

$$-14 = -a(3)$$

$$4.67 = a$$

- (e) Find the local minimum value of the function $y = x^3(x - 4)$.

Justify your answer.


$$y = x^3(x - 4)$$

$$y = x^4 - 4x^3$$


$$\frac{dy}{dx} = 4x^3 - 12x^2$$

QUESTION TWO

- (a) The diagram below shows the graph of the function $y = f(x)$.



On the axes below sketch the gradient function $y = f'(x)$.



If you need
to redraw this
graph, use the
grid on page 11.

- (b) The line $y = ax + b$ is a tangent to the graph of the function $y = 2x^2 - 3x + 1$ at the point $(3, 2)$.

ASSESSOR'S
USE ONLY

Find the values of a and b .

$$\begin{aligned}
 y &= mx + c & 2 &= 9(3) + c \\
 \frac{dy}{dx} &= 4x - 3 & 2 &= 27 + c \\
 &= 4(3) - 3 & -25 &= c \\
 &= 12 - 3 \\
 m &= 9 & a &= 9 \\
 & & b &= -25
 \end{aligned}$$

- (c) A function f is given by $f(x) = 2 - 4x + 5x^2 + ax^3$.

The gradient of the graph of the function at the point where $x = 1$ is 3.

Find the value of a .

$$\begin{aligned}
 f(x) &= 2 - 4x + 5x^2 + ax^3 \\
 f'(x) &= -4 + 10x + 3ax^2 \\
 &= -4 + 10 + 3a \\
 &= 6 + 3a \\
 -6 &= 3a \rightarrow a = -2
 \end{aligned}$$

- (d) A chemical is slowly leaking onto a floor.

The chemical spreads out from the point where it lands in a shape that can be modelled by a circle of radius r cm.

At a time t seconds after the chemical leak is noticed, r is given by $0.1t + 2$.

Use calculus to find the rate of change of area of the circle, with respect to time, when its radius is 10 cm.

(Area of circle = πr^2)

$$\begin{aligned} r &= 0.1t + 2 & (\text{area}) r' &= \frac{0.1}{2} t^2 + 2t + C \\ 10 &= 0.1t + 2 & &= \frac{0.1}{2} (80)^2 + 2(80) + C \\ 8 &= 0.1t & &= 480 + C \\ 80 &= t & &- 480 = C \end{aligned}$$

80 seconds for 10 cm radius

$$\begin{aligned} \pi (10)^2 &= 314.15 \text{ cm} \\ &= 3.14 \text{ m} \end{aligned}$$

- (e) A function is defined by $y = 3x^3 - 4a^2x + 5$ where a is a positive number.

Find the range of values of x in terms of a for which the function is decreasing.

$$\frac{dy}{dx} = 9x^2 - 8a \quad a > 0$$

QUESTION THREE

ASSESSOR'S
USE ONLY

- (a) The gradient function for a curve is given by $\frac{dy}{dx} = 3x^2 - 5$.

The curve passes through the point (1,0).

Find the equation of the curve.

$$y = \frac{3}{3}x^3 - 5x + C$$

$$y = x^3 - 5x + 4$$


$$0 = 1 - 5 + C$$

$$= -4 + C$$


$$4 = C$$

- (b) The diagram below shows the graph of the gradient function $y = f'(x)$ of a function $y = f(x)$.

ASSESSOR'S
USE ONLY



On the axes below sketch the graph of the function $y = f(x)$.



If you need
to redraw this
graph, use the
grid on page 11.

- (c) Meg is riding her motocross bike.

When she passes a fixed point P on the track, she has a speed, v , of 5 m s^{-1} , and her acceleration, a , is 0.6 m s^{-2} .

- (i) If she were to continue to accelerate at this rate, what is her speed when she has been riding for 10 seconds after passing P?

$$a = v'$$

$$0.6$$

$$v = 5$$

$$v \times 10 \text{ sec} = 50$$

- (ii) How far will she have travelled from P when she reaches a speed of 8 m s^{-1} ?

Question Three continues
on the following page.

- (iii) Meg's friend Leo was riding with her, but he begins to decelerate when they reach a speed of 8 m s^{-1} .

If he decelerates at 0.2 m s^{-2} , how far past the point P will he be when he reaches a speed of 6 m s^{-1} ?

$$a = v' = \frac{dv}{dt}$$
$$v = 8 \text{ m s}^{-1}$$

ASSESSOR'S
USE ONLY

N2

Achieved exemplar 2016

Subject:		Mathematics	Standard:	91262	Total score:	10
Q	Grade score	Annotation				
1	A3	1(b) Correct differentiation and gradient found. Not sufficient proof shown 1(c) Transfer error. Cannot allow as it made the question too simple 1(d) No correct calculus shown here 1(e) Correct differentiation but to achieve "u" grade, must be equated to zero (or implied from correct solving)				
2	M5	2(a) Parabola incorrectly placed on axes 2(b) correct solution to a and b using accurate differentiation 2(c) $a = -2$ found when the derivative is equated to zero instead of 3. 2(d) Time calculation correct, but numeric skill only 2(e) incorrect differentiation of the $4a^2 x$ term				
3	N2	3(a) Correct anti-differentiation and solving of $+c$ 3(b) wrong graph				