

No part of the candidate evidence in this exemplar material may be presented in an external assessment for the purpose of gaining credits towards an NCEA qualification.

3

91390



NEW ZEALAND QUALIFICATIONS AUTHORITY  
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD  
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

SUPERVISOR'S USE ONLY

## Level 3 Chemistry, 2017

### 91390 Demonstrate understanding of thermochemical principles and the properties of particles and substances

2.00 p.m. Wednesday 15 November 2017  
Credits: Five

Achievement	Achievement with Merit	Achievement with Excellence
Demonstrate understanding of thermochemical principles and the properties of particles and substances.	Demonstrate in-depth understanding of thermochemical principles and the properties of particles and substances.	Demonstrate comprehensive understanding of thermochemical principles and the properties of particles and substances.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

**You should attempt ALL the questions in this booklet.**

A periodic table is provided on the Resource Sheet L3–CHEMR.

If you need more room for any answer, use the extra space provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–11 in the correct order and that none of these pages is blank.

**YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.**

**Achievement**

**TOTAL**

**12**

ASSESSOR'S USE ONLY

## QUESTION ONE

ASSESSOR  
USE ONLY

(a) Complete the following table.

Symbol of particle	Electron configuration (use <i>s, p, d</i> notation)	Charge	Atomic number
Cl	$1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^1$	0	17
$Ca^{2+}$	$1s^2, 2s^2, 2p^6, 3s^2, 3p^6$	+2	20
$Mn^{2+}$	$1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2, 3d^5$	+2	25

(b) (i) Define the term electronegativity.


Electronegativity is the elements ability to attract ~~to~~ electrons. Hence to valence shell electrons.

(ii) Explain why the electronegativity of chlorine is greater than that of phosphorus.

Chlorine and Phosphorus are both in period 3. This means they have both have 3 electron shells. However Cl has 17 protons while P only has 15. Therefore Cl has more protons and the effective nuclear charge between the nucleus and valence shell electrons is greater. Thus Cl has a higher electronegativity.

- (c) The following graph shows the first ionisation energies of the Group 2 elements from Be to Ba.

ASSESSOR'S  
USE ONLY



- (i) Write an equation to show the first ionisation energy for the element calcium.



- (ii) Explain the trend shown of first ionisation energies of the Group 2 elements.

As you go down group 2 you gain an extra electron shell each time. The shielding force from the electron shell gaining is greater than the added protons effective nuclear charge. Therefore the energy required to remove <sup>one</sup> the outer ~~most~~ valence electron is less as it is easier to remove it. Thus the ionisation energy required to remove the one valence shell electron decreases as you go down group 2.

MS

## QUESTION TWO

Molecule	Boiling Point / °C	$M / \text{g mol}^{-1}$
Hydrazine, $\text{N}_2\text{H}_4$	114	32
Iodomethane, $\text{CH}_3\text{I}$	42.4	142
Decane, $\text{C}_{10}\text{H}_{22}$	174	142

Use the information in the table above to compare and contrast the boiling points of the substances below.

In your answers, you should:

- list the types of intermolecular forces present for each substance
- explain the relative strength between the particles involved.

(a) (i) Hydrazine and iodomethane.

Iodomethane has ~~no~~ temporary dipoles as its intermolecular forces. Hydrazine has temporary dipoles, permanent dipoles and hydrogen bonding for intermolecular forces. Temporary dipoles are very weak and thus even though  $\text{CH}_3\text{I}$  has a large molar mass it has a low boiling point as the energy required to break the bonds is minimal. Hydrogen bonding <sup>is</sup> ~~and~~ the strongest form of intermolecular forces and this ~~it~~ requires the most energy to break these bonds meaning  $\text{N}_2\text{H}_4$  has a high boiling point.

(ii) Iodomethane and decane.


Decane ~~has~~ and iodomethane have the same molar mass. However, decane has temporary and permanent dipoles while  $\text{CH}_3\text{I}$  just has temporary dipoles. ~~This~~ the permanent dipoles are much stronger than temporary dipoles meaning that they require more energy to break than temporary. Thus the boiling point of  $\text{C}_{10}\text{H}_{22}$  is much higher than  $\text{CH}_3\text{I}$ .

- (b) Explain why the solubility of hydrazine in water is greater than that of decane in water.

ASSESSOR'S  
USE ONLY

Solubility is determined by strength of the  $H_2O$  bond vs the intermolecular forces of the solute. As  $N_2H_4$  has hydrogen bonds (strongest permanent dipoles) it has the strongest intermolecular forces as the hydrogens become protons which then link onto the  $N_2H_4$  molecules. Decane has weaker permanent dipoles. ~~and thus~~ As  $N_2H_4$  has stronger intermolecular forces it is more likely to overcome the  $H_2O$  intermolecular forces (hydrogen bonding) and thus dissolve into  $H_2O$ .

- (c) Carbon dioxide and water are formed when decane burns completely in oxygen. The reaction is shown in the equation below.






Calculate the enthalpy of combustion for decane, given the following data:

$$\Delta_f H^\circ (C_{10}H_{22}(l)) = -301 \text{ kJ mol}^{-1}$$

$$\Delta_f H^\circ (C_{10}H_{22})$$

$$\Delta_c H^\circ (C) = -393 \text{ kJ mol}^{-1}$$


$$\Delta_c H^\circ (H_2) = -286 \text{ kJ mol}^{-1}$$



$$11 \times -286 + 10 \times -393 + 301$$

$$= -6775 \text{ kJ mol}^{-1}$$

- (d) The reaction for the complete combustion of hydrazine is shown in the equation below.



This is an exothermic reaction.

Explain the entropy changes associated with this reaction.

\* Entropy is the disorder in a system.  
 The liquid and the gas has changed to 3 gas moles. This is an increase in entropy of the system as not only have bonds been broken, there is also an increase in the amount of gaseous moles. As there is an increase in entropy in the system there is a decrease in entropy of the surroundings. This reaction will be spontaneous as  $\Delta G = \Delta H - T\Delta S$  and  $\Delta H$  is negative as the reaction is exothermic and  $\Delta S$  is positive, thus it also will subtract giving you a negative  $\Delta G$  (Gibbs free energy) and thus making it a spontaneous reaction.

A3

### QUESTION THREE

 ASSESSOR'S  
USE ONLY

Chlorine,  $\text{Cl}_2$ , bromine,  $\text{Br}_2$ , and iodine,  $\text{I}_2$ , are all halogens.  
Bromine is a liquid at room temperature.


- (a) (i) In the box below, tick the type(s) of intermolecular attractions in **liquid** bromine.

Intermolecular attraction	Tick (✓)
Temporary dipole-dipole attractions	✓
Permanent dipole-dipole attractions	
Hydrogen bonding	

- (ii) Explain why bromine is a liquid at room temperature, whereas chlorine is a gas.

Bromine is a much larger (atom) than Chlorine. Thus it takes a lot more energy to break the bonds between the <sup>liquid</sup> ~~solid~~ and <sup>gaseous</sup> ~~liquid~~ state. Therefore bromine has a higher <sup>boiling</sup> ~~melting~~ point than chlorine and it is a liquid at room temperature while chlorine is a gas. ✓

- (b) (i) Write an equation for the sublimation of iodine in the box below.



- (ii) Define the enthalpy of sublimation for iodine.

The energy required to change ~~the~~ 1 mole of  $\text{I}_2$  from the ~~gaseous~~ solid state to the gaseous state. ✓

Question Three continues  
on the following page.

$$\Delta G = \Delta H - TS$$


8

- (iii) Explain why the sublimation of iodine is spontaneous, even though the enthalpy of sublimation is a positive value.

As  $I_2$  is going from a solid to a gas, the increase in entropy in the system is very large. This increase in entropy is larger than the endothermic positive enthalpy and using the equation  $\Delta G = \Delta H - \Delta ST$  we can see if  $\Delta S$  is greater than  $\Delta H$  then  $\Delta G$  will be negative and the reaction will be spontaneous.

- (c) Iodine forms a linear  $I_3^-$  ion.

- (i) Draw the Lewis structure for the  $I_3^-$  ion in the box below.  $22 e^-$



- (ii) Explain why the  $I_3^-$  ion has a linear shape.

By 2 out of the 5 electron clouds are bonded. Thus the bond angles are  $180^\circ$  as VSEPR states that they must be as far away as possible from each other which is  $180^\circ$  and this gives  $I_3^-$  a linear shape.



(iii)  $\text{IF}_5$  has a square pyramidal shape.

ASSESSOR'S  
USE ONLY

Indicate whether the molecule  $\text{IF}_5$  is polar or non-polar.

Circle your choice.

polar

non-polar

Justify your choice.

$\text{IF}_5$  has 6 electron clouds <sup>of</sup> and which 5 of them are bonded. Due to VSEPR (valence shell <sup>electron</sup> pair repulsion) they these will as far away as possible from each other. This gives ~~the~~  $\text{IF}_5$  the square pyramidal shape. Due to F being <sup>is</sup> more electronegative than I. This means electrons will spend more time around F than I as it attracts electrons more. This results in a slightly negative charge  $\delta^-$  on the F atoms and a slightly positive charge  $\delta^+$  on the I atoms. Due to this we have <sup>polar</sup> dipoles created. As the square pyramidal shape is not symmetrical the polar dipoles will not cancel and thus  $\text{IF}_5$  is ~~indeed~~ indeed polar.

A4

# Achievement exemplar 2017

<b>Subject:</b>		<b>Chemistry</b>	<b>Standard:</b>	<b>91390</b>	<b>Total score:</b>	<b>12</b>
<b>Q</b>	<b>Grade score</b>	<b>Annotation</b>				
1	M5	In order to achieve excellence level answers, the candidate needs to acknowledge that both elements being in the third period means the same shielding effect. They also need to explain that as you go down a group, there are more protons in the nucleus, increasing nuclear charge but this is outweighed by the increasing distance between the nucleus and outer shell electrons.				
2	A3	There is some understanding of intermolecular forces but the candidate needs to be able to identify the correct forces for more than one molecule as well as compare the molecules in each example in order to demonstrate sufficient understanding. It is important to be very explicit about intermolecular forces versus forces within a molecule and ensure that solubility is understood.				
3	A4	The candidate could have referred to bromine as a molecule to demonstrate understanding of what occurs when a molecule changes state. It is important to link statements regarding the increase of entropy to what is happening to the particles for the change in entropy. When describing a shape, it is expected that the geometry of the molecule as well as the shape is communicated.				