

L3-CALCF



NEW ZEALAND QUALIFICATIONS AUTHORITY
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

Level 3 Calculus, 2017

9.30 a.m. Thursday 23 November 2017

FORMULAE AND TABLES BOOKLET for 91577, 91578 and 91579

Refer to this booklet to answer the questions in your Question and Answer booklets.

Check that this booklet has pages 2–4 in the correct order and that none of these pages is blank.

YOU MAY KEEP THIS BOOKLET AT THE END OF THE EXAMINATION.

MATHEMATICS – USEFUL FORMULAE

ALGEBRA

Quadratics

If $ax^2 + bx + c = 0$

then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Logarithms

$y = \log_b x \Leftrightarrow x = b^y$

$\log_b(xy) = \log_b x + \log_b y$

$\log_b\left(\frac{x}{y}\right) = \log_b x - \log_b y$

$\log_b(x^n) = n \log_b x$

$\log_b x = \frac{\log_a x}{\log_a b}$

Complex numbers

$z = x + iy$

$= r \operatorname{cis} \theta$

$= r(\cos \theta + i \sin \theta)$

$\bar{z} = x - iy$

$= r \operatorname{cis}(-\theta)$

$= r(\cos \theta - i \sin \theta)$

$r = |z| = \sqrt{z\bar{z}} = \sqrt{(x^2 + y^2)}$

$\theta = \arg z$

where $\cos \theta = \frac{x}{r}$

and $\sin \theta = \frac{y}{r}$

De Moivre's Theorem

If n is any integer, then

$(r \operatorname{cis} \theta)^n = r^n \operatorname{cis}(n\theta)$

COORDINATE GEOMETRY

Straight Line

Equation $y - y_1 = m(x - x_1)$

CALCULUS

Differentiation

$y = f(x)$	$\frac{dy}{dx} = f'(x)$
$\ln x$	$\frac{1}{x}$
e^{ax}	ae^{ax}
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
$\tan x$	$\sec^2 x$
$\sec x$	$\sec x \tan x$
$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$
$\cot x$	$-\operatorname{cosec}^2 x$

Integration

$f(x)$	$\int f(x) dx$
x^n	$\frac{x^{n+1}}{n+1} + c$ ($n \neq -1$)
$\frac{1}{x}$	$\ln x + c$
$\frac{f'(x)}{f(x)}$	$\ln f(x) + c$

Parametric Function

$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$

$\frac{d^2y}{dx^2} = \frac{d}{dt} \left(\frac{dy}{dx} \right) \cdot \frac{dt}{dx}$

Product Rule

$$(f \cdot g)' = f \cdot g' + g \cdot f' \quad \text{or if } y = uv \text{ then } \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

Quotient Rule

$$\left(\frac{f}{g}\right)' = \frac{g \cdot f' - f \cdot g'}{g^2} \quad \text{or if } y = \frac{u}{v} \text{ then } \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Composite Function or Chain Rule

$$(f(g))' = f'(g) \cdot g'$$

$$\text{or if } y = f(u) \text{ and } u = g(x) \text{ then } \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

NUMERICAL METHODS**Trapezium Rule**

$$\int_a^b f(x) \, dx \approx \frac{1}{2} h [y_0 + y_n + 2(y_1 + y_2 + \dots + y_{n-1})]$$

$$\text{where } h = \frac{b-a}{n} \text{ and } y_r = f(x_r)$$

Simpson's Rule

$$\int_a^b f(x) \, dx \approx \frac{1}{3} h [y_0 + y_n + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$$

$$\text{where } h = \frac{b-a}{n}, y_r = f(x_r) \text{ and } n \text{ is even.}$$


TRIGONOMETRY

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

**Sine Rule**

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Cosine Rule

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\cot^2 \theta + 1 = \operatorname{cosec}^2 \theta$$

General Solutions

$$\text{If } \sin \theta = \sin \alpha \text{ then } \theta = n\pi + (-1)^n \alpha$$

$$\text{If } \cos \theta = \cos \alpha \text{ then } \theta = 2n\pi \pm \alpha$$

$$\text{If } \tan \theta = \tan \alpha \text{ then } \theta = n\pi + \alpha$$

where n is any integer

Compound Angles

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

Double Angles

$$\sin 2A = 2 \sin A \cos A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$= 2 \cos^2 A - 1$$

$$= 1 - 2 \sin^2 A$$

Products

$$2 \sin A \cos B = \sin(A+B) + \sin(A-B)$$

$$2 \cos A \sin B = \sin(A+B) - \sin(A-B)$$

$$2 \cos A \cos B = \cos(A+B) + \cos(A-B)$$

$$2 \sin A \sin B = \cos(A-B) - \cos(A+B)$$

Sums

$$\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$$

$$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\cos C - \cos D = -2 \sin \frac{C+D}{2} \sin \frac{C-D}{2}$$

MEASUREMENT**Triangle**

$$\text{Area} = \frac{1}{2} ab \sin C$$

Trapezium

$$\text{Area} = \frac{1}{2} (a+b)h$$

Sector

$$\text{Area} = \frac{1}{2} r^2 \theta$$

$$\text{Arc length} = r\theta$$

Cylinder

$$\text{Volume} = \pi r^2 h$$

$$\text{Curved surface area} = 2\pi r h$$

Cone

$$\text{Volume} = \frac{1}{3} \pi r^2 h$$

$$\text{Curved surface area} = \pi r l \text{ where } l = \text{slant height}$$

Sphere

$$\text{Volume} = \frac{4}{3} \pi r^3$$

$$\text{Surface area} = 4\pi r^2$$