

91166



NEW ZEALAND QUALIFICATIONS AUTHORITY  
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD  
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

2

SUPERVISOR'S USE ONLY

## Level 2 Chemistry, 2018

### 91166 Demonstrate understanding of chemical reactivity

9.30 a.m. Monday 26 November 2018  
Credits: Four

| Achievement                                       | Achievement with Merit                                     | Achievement with Excellence                                     |
|---|--|---|
| Demonstrate understanding of chemical reactivity. | Demonstrate in-depth understanding of chemical reactivity. | Demonstrate comprehensive understanding of chemical reactivity. |

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

**You should attempt ALL the questions in this booklet.**

A periodic table is provided in the Resource Booklet L2–CHEMR.

If you need more room for any answer, use the extra space provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–12 in the correct order and that none of these pages is blank.

**YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.**

TOTAL

ASSESSOR'S USE ONLY



(b) Consider the following observations in another experiment using hydrogen peroxide:

- When hydrogen peroxide is mixed with solution **X**, which contains universal indicator, the colour changes from blue to green to yellow to orange-red **over a time of one hour**.
- If a crystal of **ammonium molybdate** is added to solution **X** before the hydrogen peroxide is added, the same colour changes will be seen in **three to four minutes**.

(i) Identify and explain the role of ammonium molybdate.

Use a diagram and refer to activation energy in your answer.

---

---

---

---

---

---

---

---

(ii) The pH of the original solution **X** is 10.8.

Calculate the hydronium ion concentration,  $[\text{H}_3\text{O}^+]$ , and the hydroxide ion concentration,  $[\text{OH}^-]$ , in the solution.

$[\text{H}_3\text{O}^+] =$  \_\_\_\_\_

---

---

$[\text{OH}^-] =$  \_\_\_\_\_

---

---

- (iii) The sodium hydroxide solution,  $\text{NaOH}(aq)$ , used to prepare solution **X** has a concentration of  $0.0125 \text{ mol L}^{-1}$ .

Calculate the pH of the sodium hydroxide solution.

---

---

---

---

---

---

---

---

- (iv) Another chemical in solution **X** is a salt, sodium ethanoate,  $\text{CH}_3\text{COONa}$ . When solid sodium ethanoate is dissolved in water, it separates into ions.

Use TWO relevant equations to explain whether **the solution** is acidic or basic.

---

---

---

---

---


---

---

---

**QUESTION TWO**

The Contact Process is used industrially in the manufacture of sulfuric acid. One step in this process is the oxidation of sulfur dioxide,  $\text{SO}_2(\text{g})$ , to sulfur trioxide,  $\text{SO}_3(\text{g})$ .



- (a) Write the equilibrium constant expression for this reaction.

|         |
|---------|
| $K_c =$ |
|---------|

- (b) (i) Calculate the equilibrium constant ( $K_c$ ) for this reaction at  $600^\circ\text{C}$  using the following concentrations:

$$\begin{aligned}[\text{SO}_2] &= 0.100 \text{ mol L}^{-1} \\ [\text{O}_2] &= 0.200 \text{ mol L}^{-1} \\ [\text{SO}_3] &= 0.0930 \text{ mol L}^{-1}\end{aligned}$$

---

---

---

---

- (ii) Explain what the size of the  $K_c$  value indicates about the extent of the reaction at equilibrium.

---

---

---

---



- (e) When the reaction is carried out at 450°C, the  $K_c$  value is higher than the value at 600°C.

Justify whether the oxidation of sulfur dioxide gas,  $\text{SO}_2(\text{g})$ , to sulfur trioxide gas,  $\text{SO}_3(\text{g})$ , is exothermic or endothermic.

---

---

---

---

---


---

---

---

---

---







- (c) The table below gives the pH of solutions of ethanoic acid,  $\text{CH}_3\text{COOH}(aq)$ , and nitric acid,  $\text{HNO}_3(aq)$ , of concentrations of  $0.200 \text{ mol L}^{-1}$ .

| Solution | $\text{CH}_3\text{COOH}(aq)$ | $\text{HNO}_3(aq)$ |
|----------|------------------------------|--------------------|
| pH       | 2.73                         | 0.70               |

- (i) Use the pH values to analyse the strength of the acids by calculating the concentration of their  $\text{H}_3\text{O}^+$  ions.

---

---

---

---

---

---

---

---

---

---

- (ii) Using your calculations in (i) above, predict the rate of reaction of each acid with a 2 cm strip of cleaned magnesium ribbon, Mg.

Refer to the collision theory in your answer.

---

---

---

---

---

---

---

---

---

---





