



91157

Mana Tohu Mātauranga o Aotearoa New Zealand Qualifications Authority

Level 2 Biology 2024

91157 Demonstrate understanding of genetic variation and change

Credits: Four

Achievement	Achievement with Merit	Achievement with Excellence
Demonstrate understanding of genetic variation and change.	Demonstrate in-depth understanding of genetic variation and change.	Demonstrate comprehensive understanding of genetic variation and change.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should attempt ALL the questions in this booklet.

If you need more room for any answer, use the extra space provided at the back of this booklet.

Check that this booklet has pages 2–16 in the correct order and that none of these pages is blank.

Do not write in the margins (1/1/1/2). This area will be cut off when the booklet is marked.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.

QUESTION ONE: Cats

Some domestic cats exhibit a complete dominance pattern in coat colour. The allele for black fur (B) is dominant over the allele for brown fur (b). The gene for tail length is not linked and is located on a different chromosome from the coat colour gene. The allele for long tails (L) is dominant to the allele for short tails (l).

A cat that is homozygous for both black fur and long tail is crossed with a cat that is homozygous for both brown fur and short tail.



(a) Specify the genotype of individuals in the F1 generation produced by this cross.

(b) Use the Punnett square below to show:

F

- the F1 gametes resulting from the cross and
- the possible genotypes within the F2 generation of cats.

F1 gametes

gametes			
gametes			

(c) Give the expected phenotype ratio resulting from this cross, indicating the phenotype each value represents.

3

Some cats have fur colour that is an example of co-dominance AND sex linkage. In cats with orange fur, phaeomelanin (orange pigment) completely replaces eumelanin (black or brown pigment). This gene is located on the X chromosome. The orange fur allele is (F) and is co-dominant with non-orange (f). Males can typically only be orange or non-orange (black, brown, etc.).

Sex determination in cats is the same as humans.

Female cats can have orange fur, fur without any orange (black, brown, etc.), or have tortoiseshell fur (see image on the right), in which some parts of the fur are orange and others are non-orange.

Some cat diseases are known to be sex-linked as well. Male cats have been found to be more susceptible to recessive, sex-linked diseases than female cats.



Female black tortoiseshell.

(d) Evaluate the inheritance patterns of cats to include complete dominance, co-dominance, and sex-linkage.

In your answer, refer to the examples above and include a discussion of:

- the two patterns of dominance
- why only female cats can have tortoiseshell fur colour
- the similarities and differences of recessive and dominant sex-linked genes.

Δ	
·	
	· · ·
	//
	* * *
	· · · · · · · · · · · · · · · · · · ·
	······································
Biology 91157, 2024	
	× ×

1111

5	

QUESTION TWO: Takahē

The flightless takahē (*Porphyrio hochstetteri*) has special cultural, spiritual, and traditional significance to Ngāi Tahu, the iwi from New Zealand's South Island. Ngāi Tahu value takahē as a taonga (treasure), and they continue to act as kaitiaki (guardians) of the takahē, working alongside the Department of Conservation/Te Papa Atawhai (DOC).



Takahē in a protected colony.

Genetic analyses and fossil records show that takahē were restricted to isolated areas in the north-western South Island at the height of the last ice age, approximately 29000–19000 years ago. As the climate warmed, takahē shifted their distribution, migrating to eastern and southern regions. The takahē in the north-west South Island became locally extinct. Pressures from hunting, introduced predators, habitat destruction, and competition for food led to their decline and an extreme genetic bottleneck.

After being presumed extinct for nearly 50 years, the takahē was famously rediscovered in 1948. The rediscovery of the takahē led to New Zealand's longest-running, endangered species programme. For more than 70 years, measures to protect and increase numbers of takahē have included predator control, captive breeding, and island translocations (moving small populations of birds to offshore islands).

Ongoing genetic analyses have found that introduced island populations of takahē have significantly lower levels of genetic variation than the main Fiordland population. The island population also has significantly different gene frequencies, with some alleles becoming fixed (with no variability in the gene pools) on the island sanctuaries.

Discuss the decline in genetic diversity in the takahē, with reference to the information provided.

In your answer, include discussion of:

- the terms population bottleneck, founder effect, and genetic drift
- how the genetic diversity of the gene pools of the takahē have been impacted by these processes
- why the reduced genetic diversity from island translocation is a problem for the takahē population and how this may be improved in the future.

	7	
-		
-		
		-
1		-
-		
-		-
-		
		-
		-
1		
		-
		-
-		
-		
-		
		-
1		-
		-
-		-
-		
-		-
1		
-		
		-
		-
1		_
1		_
		-
		-
1		
		_
1		
1		
]		-
		-
		_

1-1-1-

ŏ	
Biology 91157 2024	
Diology Strot, 2024	

.

This page has been deliberately left blank. The assessment continues on the following page.

9

QUESTION THREE: Lethal alleles

In 1907, Erwin Baur carried out research on the snapdragon plant, Antirrhinum majus, and studied the condition known as 'aurea', in which some plants produced golden leaves instead of green leaves. In this plant, the golden-leaf allele (G) is dominant to the green-leaf allele (q). When crossed with its own type (aurea × aurea), Baur observed a 2:1 phenotype ratio of golden:green-leafed plants, instead of the expected 3:1 ratio in the offspring.

By carrying out a number of test crosses, Baur concluded that all of the surviving golden-leafed plants were heterozygous. Homozygous dominant (GG) aurea plants lacked normal chlorophyll development and never survived.

Baur is now recognised as the first scientist to discover lethal alleles in a plant, although they had already been recognised in animals, including humans.



Describe what is meant by the term lethal allele.

(a)

(b) In the space below, show the Punnett square for a test cross between a green-leaf plant and a surviving, golden-leaf plant:



(c) This lethal allele gives a dominant, non-lethal phenotype in the heterozygote. However, we say that the lethality (ability to prevent survival) is recessive in the snapdragon, even though the colour phenotype is dominant.

Using the information provided, discuss why this snapdragon allele must be recessive for lethality and why dominant lethal alleles are rare but can be found in some adult populations, including humans.

In your answer, include discussion of:

- the terms and meaning of dominant and recessive alleles
- why the snapdragon's allele must be recessive for lethality and how the test cross shows this
- why dominant lethal alleles are rarer than recessive ones, but can sometimes exist in adult organisms.

12	
Biology 91157, 2024	

13

		14	
	Г		
		Extra space if required.	
QUESTION		Write the question number(s) if applicable.	
NUMBER			

	Extra space if required.	
QUESTION NUMBER	write the question number(s) if applicable.	

-

QUESTION	Extra space if required. Write the question number(s) if applicable.	
NOWBER		

Acknowledgements

Material from the following sources has been adapted for use in this assessment:

Page 2

https://stock.adobe.com/207291020

Page 3

https://en.wikipedia.org/wiki/Cat_coat_genetics

Page 6

https://www.doc.govt.nz/nature/native-animals/birds/birds-a-z/takahe/

Page 10 https://stock.adobe.com/324607740 ///

> > ~ ~ ~

× /+ * + * * * / · /