

No part of the candidate evidence in this exemplar material may be presented in an external assessment for the purpose of gaining credits towards an NCEA qualification.

2

91165



911650



NEW ZEALAND QUALIFICATIONS AUTHORITY  
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD  
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

SUPERVISOR'S USE ONLY

## Level 2 Chemistry, 2015

### 91165 Demonstrate understanding of the properties of selected organic compounds

9.30 a.m. Monday 23 November 2015  
Credits: Four

Achievement	Achievement with Merit	Achievement with Excellence
Demonstrate understanding of the properties of selected organic compounds.	Demonstrate in-depth understanding of the properties of selected organic compounds.	Demonstrate comprehensive understanding of the properties of selected organic compounds.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

**You should attempt ALL the questions in this booklet.**

A periodic table is provided on the Resource Sheet L2-CHEMR.

If you need more room for any answer, use the extra space provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–11 in the correct order and that none of these pages is blank.

**YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.**

Low Merit

TOTAL

16

ASSESSOR'S USE ONLY


# QUESTION ONE

- (a) (i) Complete the following table to show the structural formula and IUPAC (systematic) name for each compound.

Structural formula	IUPAC (systematic) name
	propan-1-amine
	2-chlorobutanoic acid
$\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \underset{\text{CH}_3}{\text{CH}} - \underset{\text{OH}}{\text{CH}} - \text{CH}_3$	<u>3-methylhexan-2-ol</u>
	<u>2-bromo-2-methylpropane</u> <del>2-methyl</del>

- (ii) The organic compound, 4-chloro-3-methylpent-4-ene has been named incorrectly.

Draw the implied structure and explain why it is named incorrectly.



You name from the lowest functional group number to the first ~~functional~~ group, not just from left to right

The correct IUPAC name for this structure is:

2-chloro-3-methylpent-1-ene

- (b) Butan-1-ol has the molecular formula  $C_4H_{10}O$ . Its structural formula is:



- (i) Define the term constitutional (structural) isomer.

A constitutional isomer has the same molecular formula but different structural formula.

- (ii) Draw THREE other constitutional (structural) isomers of  $C_4H_{10}O$ .

Alcohol	Structural formula
A	$  \begin{array}{ccccccc}  & H & & H & & OH & & H \\  &   & &   & &   & &   \\  H & - C & - & C & - & C & - & C - H \\  &   & &   & &   & &   \\  & H & & H & & H & & H  \end{array}  $
B	$  \begin{array}{ccccccc}  & H & & CH_3 & & OH & & H \\  &   & &   & &   & &   \\  H & - C & - & C & - & C & - & C - H \\  &   & &   & &   & &   \\  & H & & H & & H & & H  \end{array}  $
C	$  \begin{array}{ccc}  & CH_3 & \\  &   & \\  HO & - C & - CH_3 \\  &   & \\  & CH_3 &  \end{array}  $

- (iii) Choose a **secondary** alcohol from the structures above and give a reason for your choice.

Letter: A ~~B~~ ~~C~~ (circle your choice)

Reason:

The -OH group is attached to a carbon atom that is attached to two other carbons.



(c) Four separate colourless organic liquids are known to be:

- ethanol
- ethanoic acid
- hex-2-ene
- hexan-1-amine (1-aminohexane). - smells funny.

Write a procedure to identify each of these organic liquids using **only** the reagents listed below.

- acidified dichromate solution,  $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+(\text{aq})$
- bromine water,  $\text{Br}_2(\text{aq})$  - hex-2-ene - addition.
- sodium carbonate solution,  $\text{Na}_2\text{CO}_3(\text{aq})$ .

In your answer, you should:

- identify the test reagents used
- describe any observations that would be made
- identify the type of reaction that occurs
- identify the organic product of any reaction.

You do not need to include equations in your answer.

1. Smell all liquids - the one with a pungent, unpleasant and mildly fishy smell will be 1-aminohexane.

Test the remaining 3 liquids with the  $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+(\text{aq})$

The ethanoic acid and  $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+(\text{aq})$  will undergo an oxidation reaction, forming a diol.

Hex-2-ene and ~~the~~ ethanoic acid will not react and will remain green. Now test the two remaining liquids (new solutions) with bromine water. Hex-2-ene will undergo a rapid addition reaction, with the orange  $\text{Br}_2(\text{aq})$  and colourless hex-2-ene rapidly changing back to colourless. The product will


be 2,3-dibromohexane. ~~the~~ with the remaining solution, add the  $\text{Na}_2\text{CO}_3(\text{aq})$ . This should produce an acid base reaction, forming a metal salt <sup>given off</sup> carbon dioxide and water. gas/bubbles should be seen and  $\text{Na}_2\text{CO}_3$  should disappear.



→ colour changes from green to orange as the  $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+$  conc. is reduced.

## QUESTION TWO


Cling Wrap is a polymer that can be made from the monomer 1,1-dichloroethene.



1,1-dichloroethene

<http://savingcentswithcoupons.com/money-maker-deal-on-glad-cling-wrap-at-shoprite/>

- (a) (i) In the box below, draw THREE repeating units of the polymer formed.




- (ii) Explain why 1,1-dichloroethene cannot exist as a *cis-trans* isomer.

For a *cis-trans* isomer to occur, there must be a double bond (or ring) which restricts rotation and also two different groups of atoms or atoms attached to either side of the double bond. 1,1-dichloroethene does not fit this, having 2 groups of the same

- (iii) A structural isomer of 1,1-dichloroethene **can** exist as *cis-trans* isomers. Draw and name the *cis-trans* isomers.

Structure	$\begin{array}{c} \text{Cl} & & \text{H} \\ & \diagdown & / \\ & \text{C} = \text{C} \\ & / & \diagdown \\ \text{H} & & \text{Cl} \end{array}$	$\begin{array}{c} \text{H} & & \text{H} \\ & \diagdown & / \\ & \text{C} = \text{C} \\ & / & \diagdown \\ \text{Cl} & & \text{Cl} \end{array}$
Name	transdichloroethene	cisdichloroethene





In Reaction 1, propan-2-ol can be converted to propene.

In Reaction 2, propene can be converted back to propan-2-ol.

Analyse BOTH of these reactions by:


- describing the reagents and conditions needed for each reaction to occur
- identifying each type of reaction and explaining your choice
- explaining why Reaction 1 forms only a single organic product, but Reaction 2 forms a mixture of organic products.

Reaction one is an elimination reaction, where the propan-2-ol is dehydrated or eliminated using concentrated sulfuric acid to form propene. This is an elimination reaction because bonds have been broken and -OH group taken off and a double bond replacing them. There is only one product because on either side of the C the -OH is joined to, the C's both have 3 hydrogen atoms attached, so it doesn't matter if the double bond is to the right or left and by the IUPAC name would be the same. Reaction 2 is an addition reaction involving  $\text{H}_2\text{O}/\text{H}^+$  where the  $\text{H}^+$  is an addition reaction because the double bond in propene is broken and an -OH and an H group have been added. There are major and minor products available from this reaction which can be figured out using the rule 'rich get richer'. Propan-2-ol is the major product as it had the most H's attached before the bond was broken and gained another, so it got richer. Propan-1-ol would be the minor product. Reaction 2 forms two products because there are two structurally different possibilities that can occur as a product of the reaction.

E7




## QUESTION THREE



- (i) Complete the scheme above by drawing the structural formulae of the organic compounds A to D.
- (ii) Circle the functional group of each of the organic compounds A, B, and C that you have drawn.
- (iii) Identify reagents X and Y.

Reagent X: concentrated  $\text{H}_2\text{SO}_4$

Reagent Y: alcoholic  $\text{KOH}$



- (b) Ethene,  $\text{C}_2\text{H}_4(\text{g})$ , reacts with aqueous potassium permanganate solution,  $\text{KMnO}_4(\text{aq})$ , dilute acid,  $\text{H}_2\text{O}/\text{H}^+$ , and hydrogen bromide,  $\text{HBr}$ .


Compare and contrast the reactions of ethene gas with each of these three reagents.

In your answer, you should:

- describe any observations that can be made
- identify, with reasons, the type of reaction ethene undergoes with each reagent
- describe the functional group of the products formed
- include equations showing the structural formulae for the organic compounds for each reaction.

When ethene and  $\text{KMnO}_4(\text{aq})$  react, ethene undergoes an oxidation reaction, meaning the purple solution changes to a brown solution as the  $\text{MnO}_4^{2-}$  is


reduced. The Ethene forms  $\text{H}-\overset{\text{OH}}{\underset{\text{H}}{\text{C}}}-\overset{\text{H}}{\underset{\text{H}}{\text{C}}}-\text{H}$ , this is an addition reaction as the double bond in ethene has been broken and a  $-\text{OH}$  and  $-\text{H}$  group put in (2 single bonds).



Ethene will react with  $\text{H}_2\text{O}/\text{H}^+$  to form an alkane.  $\text{H}-\overset{\text{H}}{\text{C}}=\overset{\text{H}}{\text{C}}-\text{H} \xrightarrow{\text{H}_2\text{O}/\text{H}^+} \text{H}-\overset{\text{H}}{\text{C}}(\text{OH})-\overset{\text{H}}{\text{C}}(\text{H})-\text{H}$ . One product available as there are adding  $-\text{OH}/-\text{H}$  to either C would result in same structure.

This is also an addition reaction as the double bond in ethene has been broken and 2 new single bonds with  $-\text{H}$  groups added.  $\text{H}_2\text{O}/\text{H}^+$  is colourless, as is ethane, so ~~no obvious~~ observations would be made.

when ethene reacts with  $\text{HBr}$  it undergoes an addition reaction also, however as it is also breaking the double bond and replacing it with two single bonds  $\text{C}-\text{H}$  and  $\text{C}-\text{Br}$ .



This would form a haloalkane, ~~and~~ no obvious observations would occur as  $\text{HBr}$  is a colourless colour change would be from colourless to colourless, so no obvious change.

All the reactions are addition reactions, however only with  $\text{H}_2\text{O}/\text{H}^+$  and  $\text{HBr}$  are no functional groups (alkanes) are gone. With the  $\text{KMnO}_4$  no functional group is an alcohol.

Like with  $\text{KMnO}_4$ , may only one possible product and the molecule has only 2 carbons so adding  $-\text{H}$  or  $-\text{Br}$  to either would result in the same molecule, only flipped.

MS



91165



911650



NEW ZEALAND QUALIFICATIONS AUTHORITY  
 MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD  
 KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

2

SUPERVISOR'S USE ONLY

## Level 2 Chemistry, 2015

### 91165 Demonstrate understanding of the properties of selected organic compounds

9.30 a.m. Monday 23 November 2015  
 Credits: Four

Achievement	Achievement with Merit	Achievement with Excellence
Demonstrate understanding of the properties of selected organic compounds.	Demonstrate in-depth understanding of the properties of selected organic compounds.	Demonstrate comprehensive understanding of the properties of selected organic compounds.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

**You should attempt ALL the questions in this booklet.**

A periodic table is provided on the Resource Sheet L2-CHEMR.

If you need more room for any answer, use the extra space provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–11 in the correct order and that none of these pages is blank.

**YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.**

**High Merit**

**TOTAL**





**17**

ASSESSOR'S USE ONLY



(c) Four separate colourless organic liquids are known to be:

- ethanol -  $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+(\text{aq})$  orange  $\rightarrow$  green
- ethanoic acid
- hex-2-ene
- hexan-1-amine (1-aminohexane).



Write a procedure to identify each of these organic liquids using **only** the reagents listed below.

- acidified dichromate solution,  $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+(\text{aq})$
- bromine water,  $\text{Br}_2(\text{aq})$
- sodium carbonate solution,  $\text{Na}_2\text{CO}_3(\text{aq})$

In your answer, you should:

- identify the test reagents used
- describe any observations that would be made
- identify the type of reaction that occurs
- identify the organic product of any reaction.

You do not need to include equations in your answer.

First add the <sup>orange</sup>  $\text{Br}_2(\text{aq})$  water to all solutions.

This will distinguish the hex-2-ene, as the alkene will <sup>be the only one to</sup> react with the <sup>orange</sup> bromine water ( $\text{Br}_2(\text{aq})$ ) through a substitution reaction to form colourless  $\text{CH}_3\text{CHBrCHBrCH}_2\text{CH}_2\text{CH}_3$ .

This ~~reaction~~ does not require UV light or heat and is a fast reaction which occurs immediately. The product formed is a halalkane bearing 2 <sup>Br</sup> functional groups.

Next add the <sup>and heat</sup> acidified dichromate solution  $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+(\text{aq})$  to all <sup>other</sup> solutions. This will

distinguish ethanol, the alcohol as it will be the only one which reacts with the acidified dichromate <sup>with heat</sup> to form the <sup>functional</sup> carboxylic acid-ethanoic acid bearing the  $\text{COOH}$  group. The ~~ethanol~~ ethanol will go from the orange ~~dichromate~~ to green and it is


an elimination reaction, forming  $\text{H}_2\text{Cg}$  also.

~~The~~ Next add the Sodium carbonate solution ( $\text{Na}_2\text{CO}_3$ ) to the remaining 2 solutions. This distinguishes the carboxylic acid  $\text{CH}_3\text{COOH}$ , which will ~~form form~~ go through the acid + ~~base~~ carbonate  $\rightarrow$  salt + water + carbon dioxide reaction.

~~Upon~~ Upon the formation of the ~~carbon~~ Carbon dioxide ( $\text{CO}_2\text{g}$ ) gas, ~~fizzing~~ bubbles will be produced will be observed and ~~as~~  $\text{CO}_2\text{g}$  is produced. That leaves the amine which is the last solution. //

## QUESTION TWO


Cling Wrap is a polymer that can be made from the monomer 1,1-dichloroethene.



1,1-dichloroethene

<http://savingcentswithcoupons.com/money-maker-deal-on-glad-cling-wrap-at-shoprite/>

- (a) (i) In the box below, draw THREE repeating units of the polymer formed.




- (ii) Explain why 1,1-dichloroethene cannot exist as a *cis-trans* isomer.

Because geometric isomers can only exist when ~~the~~ <sup>same or different</sup> atoms or group of atoms are on ~~the~~ <sup>each</sup> ~~carbon~~ <sup>carbon participating</sup> bond ~~is~~ <sup>are</sup> in the double.





- (iii) A structural isomer of 1,1-dichloroethene can exist as *cis-trans* isomers.

Draw and name the *cis-trans* isomers.



Structure	$\begin{array}{c} \text{Cl} \quad \text{Cl} \\ \diagdown \quad \diagup \\ \text{C} = \text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$	$\begin{array}{c} \text{Cl} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C} = \text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{Cl} \end{array}$
Name	<u>cis-1,2-dichloroethene</u>	<u>trans-1,2-dichloroethene</u>





In Reaction 1, propan-2-ol can be converted to propene.

In Reaction 2, propene can be converted back to propan-2-ol.

Analyse BOTH of these reactions by:

- describing the reagents and conditions needed for each reaction to occur
- identifying each type of reaction and explaining your choice
- explaining why Reaction 1 forms only a single organic product, but Reaction 2 forms a mixture of organic products.

Reaction 1 <sup>dehydration</sup> elimination reaction. It needs concentrated  $\text{H}_2\text{SO}_4/\text{H}^+$  and heat to occur.

The conc.  $\text{H}_2\text{SO}_4/\text{H}^+$  is the dehydrating agent and removes an OH and H from the 2 Carbon atoms to form  $\text{H}_2\text{O}$  and propene from propan-2-ol.

~~Reaction 2~~

Reaction 2 This is an addition reaction.

The  $\text{C}=\text{C}$  double bond breaks and ~~each~~ <sup>one</sup> Carbon accepts the  $-\text{OH}$  group and the other Carbon the Hydrogen atom. They react with  $\text{H}_2\text{O}/\text{H}^+$  and heat for this to occur. This reaction can ~~accept~~ <sup>form</sup>


2 organic products, a major product (most of the products form this) and a minor product.

~~The~~ Following Markovnikov's rule "the rich get richer" So the Carbon atom bearing the most H atoms will get the ~~rest~~ <sup>other</sup> H atom from the reaction and the Carbon participating in the double bond the OH group. This is the major product. The minor product is when the Carbon bearing the most H atoms gets the OH group instead and the other C atom from the double bond gets the H atom.

See extra paper  
seen

MG

### QUESTION THREE



- Complete the scheme above by drawing the structural formulae of the organic compounds A to D.
- Circle the functional group of each of the organic compounds A, B, and C that you have drawn.
- Identify reagents X and Y.

Reagent X:  $\text{C. H}_2\text{SO}_4 / \text{H}^+$  heat


Reagent Y: \_\_\_\_\_

- (b) Ethene,  $\text{C}_2\text{H}_4(\text{g})$ , reacts with aqueous potassium permanganate solution,  $\text{KMnO}_4(\text{aq})$ , dilute acid,  $\text{H}_2\text{O}/\text{H}^+$ , and hydrogen bromide,  $\text{HBr}$ .



Compare and contrast the reactions of ethene gas with each of these three reagents.

In your answer, you should:

- describe any observations that can be made
- identify, with reasons, the type of reaction ethene undergoes with each reagent
- describe the functional group of the products formed
- include equations showing the structural formulae for the organic compounds for each reaction.







① The addition of ~~the~~ ethene with  $\text{H}_2\text{O}/\text{H}^+$  results in the ~~breaking of the double bond and the~~ addition of an -OH and H atom. ~~the reactant~~ <sup>originally involved in the double bond.</sup>

Solution remains colourless. This is an addition reaction as ~~for~~ groups are added to the original structure. The functional group is an alcohol.

② This is an ~~addition~~ <sup>oxidation</sup> reaction and ~~is~~ <sup>requires</sup> the addition of 2 -OH groups to ~~the~~ <sup>original</sup> ethene. Each Carbon atom now gains an OH group and the double bond breaks. The  $\text{KMnO}_4(\text{aq})$  is originally purple in colour ~~but~~ but the reaction will turn the solution <sup>due to presence of</sup> very pale pink, nearly colourless ( $\text{Mn}^{2+}$  ion). ~~with~~ The functional group of the -OH is alcohol.

③ The ~~addition~~ addition reaction of the H atom and Br atom. ~~The  $\text{C}=\text{C}$  bond with the  $\text{HBr}$  molecule~~ <sup>from the substance</sup> ~~breaks and this forms~~ <sup>the</sup> the carbon ~~bond~~ <sup>substance</sup> breaks and ~~this forms~~ has to carbon <sup>(C=C) double</sup> the addition of a hydrogen (H) atom and Br (Bromine) atom. (with the addition of the haloalkane -X group (-Br)). the solution goes from colourless to orange due to the <sup>presence</sup> Br atom. The functional group is haloalkane. The Br is the functional group.

M6



Extra paper if required.

Write the question number(s) if applicable.

QUESTION  
NUMBERASSESSOR  
USE ONLY

2 b) ~~This~~ Reaction 1 cannot form a major or minor product, only 1 product as Markovnikov's rule applies only with the addition of a  $\text{H-X}$  group or  $\text{-OH}$  group whereas this is an elimination reaction only 1 product can form. //