

No part of the candidate evidence in this exemplar material may be presented in an external assessment for the purpose of gaining credits towards an NCEA qualification.

2

91164



911640



NEW ZEALAND QUALIFICATIONS AUTHORITY
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

SUPERVISOR'S USE ONLY

Level 2 Chemistry, 2016

91164 Demonstrate understanding of bonding, structure, properties and energy changes

9.30 a.m. Monday 21 November 2016
Credits: Five

Achievement	Achievement with Merit	Achievement with Excellence
Demonstrate understanding of bonding, structure, properties and energy changes.	Demonstrate in-depth understanding of bonding, structure, properties and energy changes.	Demonstrate comprehensive understanding of bonding, structure, properties and energy changes.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should attempt ALL the questions in this booklet.

A periodic table is provided on the Resource Sheet L2-CHEMR.

If you need more room for any answer, use the extra space provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–10 in the correct order and that none of these pages is blank.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.

Merit

TOTAL

15

ASSESSOR'S USE ONLY

QUESTION ONE

- (a) Instant cold packs are useful for treating sports injuries on the field. They contain salts such as ammonium nitrate, NH_4NO_3 . When the packs are activated, the salt dissolves in water, causing the temperature to decrease.

Circle the term that best describes the dissolving process.

endothermic

exothermic

Give a reason for your choice.

This process is endothermic due to dissolving occurring which requires heat hence heat is absorbed (endog) and temperature decreased of surrounding due to heat being absorbed (endog)

- (b) The equation for hydrating anhydrous copper sulfate is as follows:



Circle the term that best describes this reaction.

endothermic

exothermic

Give a reason for your choice.

This reaction is exothermic due to $\Delta_r H^\circ = -78.2 \text{ kJ mol}^{-1}$ which is negative meaning heat energy is released and therefore exothermic


- (c) Pentane, C_5H_{12} , is a liquid at room temperature. It evaporates at 36.1°C in an endothermic process.

- (i) Explain why the evaporation of pentane is an endothermic process.


In order for C_5H_{12} to change state from liquid to gas to evaporate, heat energy is required to break bonds or change state hence heat is absorbed resulting in this process to be endothermic

- (ii) Draw, including labels, the energy diagram for the combustion of pentane, $C_5H_{12}(l)$.

Pentane combustion:



Include in your diagram the reactants, products, and change in enthalpy.



- (iii) Hexane, C_6H_{14} , like pentane, will combust (burn) in sufficient oxygen to produce carbon dioxide gas and water.

Hexane combustion:



Justify which alkane – pentane or hexane – will produce more heat energy when 125 g of each fuel is combusted in sufficient oxygen.

$$M(C_5H_{12}) = 72.0 \text{ g mol}^{-1}$$

$$M(C_6H_{14}) = 86.0 \text{ g mol}^{-1}$$

$$n(C_6H_{14}) = \frac{m}{M} = \frac{125}{86} = 1.453488372 \text{ moles}$$

$$2 \text{ moles } (C_6H_{14}) = 8316 \text{ kJ} \quad 1 \text{ mole} = 4158 \text{ kJ}$$

$$1.453488372 \text{ moles} = \underline{6043.604651 \text{ kJ}}$$

$$n(C_5H_{12}) = \frac{m}{M} = \frac{125}{72} = 1.736111111 \text{ moles}$$

$$2 \text{ moles} = 8316 \quad 1 \text{ mole} = 4158 \text{ kJ}$$

$$1.736111111 \text{ moles} = \underline{7218.75 \text{ kJ}}$$

∴ pentane produces more heat energy

(7218.75 kJ) than hexane which produces 6043.605 kJ. ~~Thus~~ Pentane has 1175.145 kJ more heat than hexane.

QUESTION TWO

- (a) Complete the table below by stating the type of substance, the type of particle, and the attractive forces between the particles in the solid for each substance.

Substance	Type of substance	Type of particle	Attractive forces between particles
$\text{ZnCl}_2(s)$ (zinc chloride)	<u>ionic</u>	<u>ions</u>	<u>strong ionic bonds</u>
$\text{C}(s)$ (graphite)	<u>covalent</u>	<u>atoms</u>	<u>weak covalent bonds</u>
$\text{CO}_2(s)$ (carbon dioxide/dry ice)	<u>molecular</u>	<u>molecules</u>	<u>weak intermolecular forces</u>

- (b) Carbon (graphite) conducts electricity when it is solid, whereas zinc chloride, ZnCl_2 , will not conduct electricity when solid, but will conduct when molten.

Justify this statement in terms of the particles, structure, and bonding for both substances.

Graphite is a covalent bond in a 2D lattice structure where each C atom is covalently bonded to 3 other C atoms. Forces are weak. In order to conduct electricity, the substance requires free moving charged particles which graphite has in solid state. Due to having free moving charged particles in solid state as there is a delocalized electron in graphite, graphite can conduct electricity in solid state.

Whereas ZnCl_2 is an ionic solid held in a 3D lattice structure consisting of oppositely charged ions. In order to conduct electricity, ZnCl_2 requires free moving charged particles which it does have but is held strongly into place with strong ionic bonding in solid state and therefore cannot conduct electricity. Though when ZnCl_2 is dissolved in water or is in molten state, these charged particles are no longer held in place with strong ionic bonding resulting in free moving charged particles to be present and therefore can conduct electricity in molten state but not

- (c) Solid zinc chloride, $\text{ZnCl}_2(s)$, is soluble in water. Dry ice, $\text{CO}_2(s)$, is not readily soluble in water.


ASSESSOR'S
USE ONLY

Justify these statements in terms of the particles, structure, and bonding of these substances.

You may include a diagram or diagrams in your answer.

ZnCl_2 is an ionic solid in a 3D lattice structure consisting of oppositely charged ions with strong ionic bonding. ZnCl_2 is soluble in water as despite there being strong bonding between water molecules, the water molecules are more attracted ^{to the} positive (δ^+) + negative (δ^-) ions held in ZnCl_2 causing ~~it~~ ^{$\text{Zn}^{2+} 3 \text{Cl}^-$} it to break up into ions meaning ZnCl_2 is soluble in water. Whereas CO_2 is a molecular substance made up of non polar molecules. Non polar substances are only soluble in non polar solvents which means water which is polar cannot dissolve CO_2 . This is because the attraction between water is stronger than the attraction between the ~~water and~~ CO_2 and water causing ~~the~~ CO_2 to not mix and not be soluble in water.


Space for diagrams






~~not~~

M5

QUESTION THREE

ASSESSOR'S
USE ONLY

- (a) (i) Draw the Lewis structure (electron dot diagram) for each of the following molecules, and name their shapes.

Molecule	H ₂ O	CS ₂	PH ₃
Lewis structure			
Name of shape	<u>bent</u>	<u>linear</u>	<u>triangular pyramid</u>
Approximate bond angle around the central atom	109.5°	180°	109.5°

- (ii) Compare and contrast the shapes and bond angles of H₂O, CS₂ and PH₃.

H₂O has 4 regions of electron clouds around the central O giving it a tetrahedral arrangement.

~~It has 2 bonding & 2 non bonding pairs and all 4 regions repel each other to achieve maximum stability & minimum repulsion.~~ It has 2 bonding & 2 non bonding pairs and all 4 regions repel each other to achieve maximum stability & minimum repulsion.

Non bonding pairs repel bonding pairs more than the bonding pairs repel each other as it's held closer to the nucleus. The non bonding pair contributes to the shape but is not considered part of it. This gives the shape a bent shape with an approximate angle of 109.5°.

CS₂ has 2 regions of electron clouds around the central C atom giving it a linear arrangement. All 2 regions repel each other to achieve maximum stability & minimum repulsion.

This gives it a bent shape at an approximate angle of 180°.

PH₃ has 4 regions of electron clouds similar to H₂O giving it a tetrahedral arrangement. It has 3 bonding & 1 non bonding pairs which all repel each other for maximum stability & minimum

repulsion. Non bonding pairs repel bonding pairs more than bonding pairs repel each other as they're held closer to the nucleus. Non bonding pairs contribute to the shape but are not considered part of it. This gives it a triangular pyramid shape with an

(b) The Lewis structures for two molecules are shown.

ASSESSOR'S
USE ONLY

Molecule	$\begin{array}{c} \text{H}-\ddot{\text{N}}-\text{H} \\ \\ \text{H} \end{array}$ <p>Ammonia</p>	$\begin{array}{c} \text{H}-\text{B}-\text{H} \\ \\ \text{H} \end{array}$ <p>Borane</p>
Polarity of molecule	polar	non-polar

Ammonia, NH_3 , is polar, and borane, BH_3 , is non-polar.


Justify this statement.

NH_3 has 3 bonding + 1 non-bonding pair around its central N atom giving it a trigonal pyramidal shape of an approximate angle of 107° . NH_3 has polar bonds due to the difference in electronegativity where N (δ^-) is more electronegative than H (δ^+) causing unequal sharing of electrons and net dipoles. The overall molecule is also polar due to the shape of trigonal pyramidal being asymmetrical causing net dipoles to not cancel out leaving a polar molecule. //

BH_3 has 3 bonding and 0 non-bonding pairs around the central B atom giving it a triangular planar shape of an approximate angle of 120° . BH_3 has polar bonds due to the difference in electronegativity where B (δ^-) is more electronegative than H (δ^+) causing unequal sharing of electrons and net dipoles. Due to the shape of triangular planar being symmetrical, the net dipoles can cancel each other out leaving a non-polar molecule overall. //


- (c) Calculate the enthalpy change, $\Delta_r H^\circ$, for the reaction of but-1-ene gas, $C_4H_8(g)$, with hydrogen gas, $H_2(g)$, to form butane gas, $C_4H_{10}(g)$.

Use the average bond enthalpies given in the table below.



Bond	Average bond enthalpy / kJ mol^{-1}
C=C	614
C-C	346
C-H	414
H-H	436

Show your working and include appropriate units in your answer.



Annotated Exemplar

Merit exemplar 2016

Subject:		Chemistry	Standard:	91164	Total score:	15
Q	Grade score	Annotation				
1	M5	<p>Both parts (a) and (b) are correct.</p> <p>In part (c)(i), the response outlines that energy is absorbed, but the type of bonds (intermolecular) is missing.</p> <p>In part (c)(ii), the ΔH label is missing, however, the reactants and products are correctly labelled.</p> <p>The calculation for hexane is correct, but the pentane calculation is not.</p>				
2	M5	<p>Two rows and two columns are correct in the table in part (a).</p> <p>For part (b), the response outlines that charged particles are needed to conduct electricity, but for molten zinc chloride, the charged particles are not called ions or cat/anions.</p> <p>For part (c), the relative strengths are mentioned for CO_2, although expressed the opposite way round, as well as the polarity of water. No link to polarity of water attractions to ions in zinc chloride is made.</p>				
3	M5	<p>Table completely correct in part (a)(i).</p> <p>The candidate just about explains the shapes and angles in part (b), but did not answer the question which asked for a comparison and a contrast.</p> <p>In part (b), the candidate states incorrectly that the bonds cancel, instead of dipoles or polarities cancelling.</p> <p>The candidate has included a partially correct process for the calculation, but including H—H bond under bond making, is a major error.</p>				