

Assessment Schedule – 2016**Chemistry: Demonstrate understanding of the properties of selected organic compounds (91165)****Evidence Statement**

Q	Evidence	Achievement	Merit	Excellence
ONE (a) (i)	2-iodohexane $\text{CH}_3 - \text{CH}_2 - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_2 - \underset{\text{O}}{\overset{\text{ }}{\text{C}}} - \text{OH}$ $\text{H} - \text{C} \equiv \text{C} - \text{CH}_2 - \text{CH}_3$ propan-1-amine (1-propanamine)	<ul style="list-style-type: none"> Any FOUR structures or names correct. 	<ul style="list-style-type: none"> Any NINE structures and names correct. 	
(ii)	$\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$ pentane $\text{CH}_3 - \text{CH}_2 - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_3$ methylbutane (or accept 2-methylbutane) $\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{CH}_3 \\ \\ \text{CH}_3 \end{array}$ dimethylpropane (or accept 2,2-dimethylpropane)	<ul style="list-style-type: none"> All THREE classifications correct. 		
(b)(i)	A. Tertiary (or 3°) B. Primary (or 1°) C. Secondary (or 2°)	<ul style="list-style-type: none"> All THREE classifications correct. 		
(ii)	Classifying haloalkanes as primary, secondary, or tertiary requires counting the number of C or H atoms bonded to the C atom to which the halogen is attached. If the C atom bonded to the halogen has 3 other carbon atoms (or 0 H atoms) bonded to it, the haloalkane is a tertiary (3°) alkane.	<ul style="list-style-type: none"> States how to classify haloalkanes correctly. 	<ul style="list-style-type: none"> Links how to classify haloalkanes correctly to structure A in the table. 	

<p>(c)(i)</p> <p>(ii)</p>	<p>A. <i>trans</i> B. <i>cis</i></p> <p>1,2-dibromoethene can form <i>cis</i> and <i>trans</i> isomers because it has a double bond. The double bond between two carbon atoms does not allow any rotation of atoms around it. As well as the double bond, the C atoms directly attached to it must have two different atoms or groups attached to them. For 1,2-dibromoethene, both the C atoms on the double bond have an H and a Br atom bonded to them.</p> <p>When these two requirements are met, two alkenes can have the same molecular formula and the same sequence of atoms in the formula, but a different arrangement in space (a different 3D formula), hence they are <i>cis</i> and <i>trans</i> isomers.</p>	<ul style="list-style-type: none"> • Correctly identifies both <i>cis</i> and <i>trans</i> isomers, <p>AND</p> <p>EITHER</p> <p>States that a rigid double bond is needed.</p> <p>OR</p> <p>Each carbon around the double bond needs TWO different atoms or groups attached to it.</p>	<ul style="list-style-type: none"> • Explains why the double bond is required for <i>cis</i> and <i>trans</i> isomerism. <p>OR</p> <p>Explains why each C atom on the double bond must have two different atoms or groups attached to them.</p>	<ul style="list-style-type: none"> • Explains why the double bond and two different atoms or groups of atoms on the C atoms of the double bond are required for <i>cis</i> and <i>trans</i> isomers. <p>AND</p> <p>Relates their answer specifically to the example given in the question.</p>
---------------------------	--	---	--	---

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	1a	2a	3a	4a	2m	3m	1e OR 1e (with minor error or omission) + 1m	1e + 1m

Q	Evidence	Achievement	Merit	Excellence
TWO (a)(i) (ii) (b)(i) (ii) (c)	<p>The boiling points of both alkanes and alcohols increase as the number of C atoms increases. The boiling points of alcohols are always higher than the alkanes (with the same number of C atoms).</p> <p>Alkanes with 1, 2, 3, and 4 C atoms (methane, ethane, propane, and butane) will be gases at room temperature.</p> <p>$\text{CH}_3\text{CH}_2\text{NH}_2(\text{aq}) + \text{HCl}(\text{aq}) \rightarrow \text{CH}_3\text{CH}_2\text{NH}_3^+(\text{aq}) + \text{Cl}^-(\text{aq})$ OR $\text{CH}_3\text{CH}_2\text{NH}_3\text{Cl}(\text{aq})$</p> <p>Carboxylic acids have acidic properties because when they react, some of the acid molecules donate H^+ to water molecules. $\text{CH}_3\text{COOH}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{CH}_3\text{COO}^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$</p> <p>Ethane is an alkane which will react slowly with bromine water, $\text{Br}_2(\text{aq})$, if (UV) light (or heat) is present, whereas the reaction between ethene and bromine water is immediate, and requires no special conditions. When they react, both ethane and ethene cause orange / yellow bromine water to decolourise. Since ethane is already a saturated molecule / hydrocarbon, it undergoes substitution with bromine water, resulting in the formation of bromoethane, $\text{CH}_3\text{CH}_2\text{Br}$. Ethene, however, is an unsaturated molecule / hydrocarbon, because it has a double $\text{C}=\text{C}$ bond, which breaks, allowing two extra atoms to bond to the structure, resulting in the formation of 1,2-dibromoethane, $\text{CH}_2\text{BrCH}_2\text{Br}$. This is an addition reaction.</p>	<ul style="list-style-type: none"> Correctly identifies one trend. OR Correctly identifies all four gaseous alkanes. Completes a balanced equation correctly. Identifies H^+ is transferred from ethanoic acid to water molecules. OR Identifies $\text{H}^+ / \text{H}_3\text{O}^+$ as a product of the reaction. OR Identifies an acidic property, e.g. indicator, neutralisation reaction. Identifies conditions and observations for one reaction. OR Both structures correct. Identifies both types of reaction correctly. 	<ul style="list-style-type: none"> Both trends identified. AND All four gases named. Explains why carboxylic acids have acidic properties, e.g. donates a proton to water / produces hydronium ions. AND Writes a balanced equation for a reaction. Links conditions, observations, reaction type, and structural formula of product for one or both reactions. (Does not 'compare and contrast'.) 	<ul style="list-style-type: none"> Compares and contrasts both reactions comprehensively.

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	1a	2a	3a	4a	2m	3m	1e (minor error/omission)	1e

Q	Evidence	Achievement	Merit	Excellence
THREE (a)(i) (ii) (b)(i) (ii)	<p>Compound A. $\text{CH}_3\text{—CH}_3$ Compound B. $\text{CH}_3\text{—CH}_2\text{OH}$ Compound C. $\text{CH}_3\text{—COOH}$ or $\text{CH}_3\text{—COH}$ (aldehyde) Reagent X. concentrated H_2SO_4</p> <p>Reaction 1: substitution Reaction 2: substitution Reaction 3: oxidation</p> <p> $\begin{array}{c} \text{H} \quad \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C} = \text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \quad \text{C}_6\text{H}_5 \end{array}$ </p> <p>Since the monomer for this reaction, styrene, is an alkene, when polymerisation occurs, the double bond in each styrene molecule is broken, freeing up a bonding space on each of the C atoms that was part of the double bond. This allows the monomers to join together by forming covalent bonds to make polystyrene.</p> <p>Since double bonds in styrene are being broken and molecules added into the freed up bonding spaces to make polystyrene, this is an addition reaction.</p> <p>Polymerisation reactions occur when many monomers are chemically joined.</p>	<ul style="list-style-type: none"> Any THREE correct in (a)(i). OR All THREE correct in (a)(ii). Correct monomer is drawn. Correctly states why this is an addition reaction. 	<ul style="list-style-type: none"> Any SIX correct. Links how the polymer forms to why it is an addition reaction. 	

(c)(i)	<p>Major product: $\text{CH}_3 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_3$ or 2-chloropropane</p> <p>Minor product: $\text{Cl} - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$ or 1-chloropropane</p>	<ul style="list-style-type: none"> Both major and minor products are correct (indicated by either structure or name). OR States the definition of Markovnikov's rule (NOT just the 'rich get richer'). OR Identifies propene as an asymmetric alkene. 	<ul style="list-style-type: none"> Explains why there are two products formed in this reaction. OR Explains why 2-chloropropane is the major product. OR Explains why 1-chloropropane is the minor product. 	<p>Elaborates on the reaction by explaining why two products form and how the major and minor products are determined.</p>
(ii)	<p>Two products are formed in this reaction because propene is an asymmetric alkene. When another asymmetric molecule such as hydrogen chloride, HCl, is added to it, there are two possible products. One product is produced in greater quantities (the major product) than the other (minor product).</p> <p>The rule for determining which is the major product (called the Markovnikov's rule) states that the C in the double bond with the most H atoms directly attached to it is most likely to gain another H atom ('rich get richer' concept).</p> <p>The most common product, the major product, is therefore 2-chloropropane, and 1-chloropropane is the minor product.</p> <p><i>(The term Markovnikov's rule is not required.)</i></p>			

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	1a	2a	3a	4a	2m	3m	1e OR 1e (with minor error or omission) + 1m	1e + 1m

Cut Scores

Not Achieved	Achievement	Achievement with Merit	Achievement with Excellence
0 – 7	8 – 14	15 – 18	19 – 24