


Assessment Schedule – 2021

Calculus: Apply differentiation methods in solving problems (91578)

Evidence Statement

	Expected coverage	Achievement (u)	Merit (r)	Excellence (t)
ONE (a)	$\frac{dy}{dx} = 3e^{3x} \sin(2x) + e^{3x} \cos(2x) \cdot 2$	Correct derivative.		
(b)(i) (ii)	(1) $x < 2, x = 4$ (2) $3 < x < 6$ 3	2 out of 3 correct responses.		
(c)	$y = (2x + 3)e^{x^2}$ $\frac{dy}{dx} = 2e^{x^2} + (2x + 3)(2x)e^{x^2}$ $\frac{dy}{dx} = 2e^{x^2} (1 + x(2x + 3))$ $\frac{dy}{dx} = 2e^{x^2} (2x^2 + 3x + 1)$ $\frac{dy}{dx} = 0$ for stationary points. $2e^{x^2} = 0$ has no solutions since $2e^{x^2} > 0$ $2x^2 + 3x + 1 = 0$ $x = -\frac{1}{2}$ or $x = -1$	Correct derivative.	Correct solution with correct derivative.	
(d)	$x = t^2 + 3t$ $\frac{dx}{dt} = 2t + 3$ $y = t^2 \ln(2t - 3)$ $\frac{dy}{dt} = 2t \ln(2t - 3) + \frac{2t^2}{2t - 3}$ $\frac{dy}{dx} = \frac{2t \ln(2t - 3) + \frac{2t^2}{2t - 3}}{2t + 3}$ At (10,0): $t^2 + 3t = 10$ $t^2 + 3t - 10 = 0$ $(t + 5)(t - 2) = 0$ $t = -5$ or $t = 2$ Since $t > \frac{3}{2}$, $t = 2$ $\frac{dy}{dx} = \frac{4 \ln(1) + 8}{7}$ $\frac{dy}{dx} = \frac{8}{7}$	$\frac{dy}{dt}$ correct.	$\frac{dy}{dt}$ correct And $t^2 + 3t = 10$ solved to find $t = -5$ or $t = 2$	T1: Correct solution with correct $\frac{dy}{dx}$.

<p>(e)</p>	$V = \pi r^2 h$ $= \pi r^2 (3 - 2r)$ $= 3\pi r^2 - 2\pi r^3$ $\frac{dV}{dr} = 6\pi r - 6\pi r^2$ <p>At maximum, $\frac{dV}{dr} = 0$</p> $6\pi r(1 - r) = 0$ $r = 0 \text{ (no)} \therefore r = 1$ $V = \pi 1^2 (3 - 2 \times 1) = \pi$ $\frac{d^2V}{dr^2} = 6\pi - 12\pi r$ <p>When $r = 1$, $\frac{d^2V}{dr^2} = -6\pi < 0$</p> <p>Therefore $V = \pi$ is maximum volume.</p>	 <p>Correct expression for $\frac{dV}{dr}$.</p>	<p>Correct expression for $\frac{dV}{dr}$ and finds $r = 1$.</p>	<p>T1: Correct expression for $\frac{dV}{dr}$ and shows that $V = \pi$ but does not prove it is the maximum volume with either the first or second derivative test.</p> <p>T2: Correct expression for $\frac{dV}{dr}$ and correct proof.</p>
------------	---	---	--	---

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	ONE partial solution.	1u	2u	3u	1r	2r	T1	T2 or two T1

	Expected coverage	Achievement (u)	Merit (r)	Excellence (t)
TWO (a)	$\frac{dy}{dx} = 5(1-x^2)^4 \times (-2x)$	Correct derivative.		
(b)	$\frac{dy}{dx} = \frac{(x+1)2x - x^2}{(x+1)^2}$ $= \frac{x^2 + 2x}{(x+1)^2}$ $\frac{dy}{dx} = 0 \Rightarrow x(x+2) = 0$ $x = 0 \text{ or } x = -2$	Correct solutions with correct derivative.		
(c)	$y = (x^2 + 3x + 2)\cos 3x$ $\frac{dy}{dx} = (2x + 3)\cos 3x - (x^2 + 3x + 2)3\sin 3x$ <p>Crosses y-axis $\Rightarrow x = 0, y = 2, \frac{dy}{dx} = 3$</p> <p>Normal gradient is $\frac{-1}{3}$</p> <p>Equation of normal:</p> $y - 2 = \frac{-1}{3}(x - 0)$ $y = \frac{-1}{3}x + 2$ $3y + x - 6 = 0$	Correct derivative.	Correct solution with correct derivative.	
(d)	$\frac{dV}{dt} = 60$ $V = \frac{4}{3}\pi r^3$ $\frac{dV}{dr} = 4\pi r^2$ $\frac{dr}{dt} = \frac{dV}{dt} \times \frac{dr}{dV}$ $= \frac{60}{4\pi r^2}$ $= \frac{15}{\pi r^2}$ $r = 15 \Rightarrow \frac{dr}{dt} = \frac{15}{\pi 15^2}$ $= \frac{1}{15\pi} (= 0.0212) \text{ cm s}^{-1}$	Correct expression for $\frac{dr}{dt}$.	Correct solution with correct $\frac{dr}{dt}$.	

<p>(e)</p> $y = \sqrt{2x-4}$ $\frac{dy}{dx} = \frac{1}{\sqrt{2x-4}}$ <p>Gradient of tangent = $\frac{y-1}{x+2}$</p> $\frac{1}{\sqrt{2x-4}} = \frac{\sqrt{2x-4}-1}{x+2}$ $x+2 = 2x-4 - \sqrt{2x-4}$ $\sqrt{2x-4} = x-6$ $2x-4 = x^2 - 12x + 36$ $x^2 - 14x + 40 = 0$ $(x-4)(x-10) = 0$ <p>$x = 4$ or $x = 10$</p> <p>Rejecting $x = 4$ by checking the surd equation</p> <p>$x = 10$ $\sqrt{16} = 4$ True</p> <p>$x = 4$ $\sqrt{4} = -2$ False</p> <p>One solution: $x = 10$</p> <p>Therefore, the coordinates of point P are (10,4)</p> <p>OR</p> <p>Rejecting $x = 4$ by checking the gradient:</p> <p>At (10,4), $\frac{dy}{dx} = \frac{1}{\sqrt{16}} = \frac{1}{4}$</p> <p>Gradient: $\frac{y-1}{x+2} = \frac{3}{12} = \frac{1}{4}$</p> <p>At (4,2), $\frac{dy}{dx} = \frac{1}{\sqrt{4}} = \frac{1}{2}$</p> <p>Gradient: $\frac{y-1}{x+2} = \frac{1}{6}$</p> <p>One solution: $x = 10$</p> <p>Therefore, the coordinates of point P are (10,4)</p>	<p>Correct derivative:</p> $\frac{dy}{dx} = \frac{1}{\sqrt{2x-4}}$	<p>Correct derivative:</p> $\frac{dy}{dx} = \frac{1}{\sqrt{2x-4}}$ <p>and</p> $\sqrt{2x-4} = x-6$	<p>T1:</p> <p>Correct solution with correct derivative: P (10,4) without any justification for $x \neq 4$</p> <p>T2:</p> <p>Correct solution with correct derivative: P (10,4) $x \neq 4$ must be justified with respect to either the surd equation or the gradient of the tangent.</p>
--	--	---	--

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	ONE partial solution.	1u	2u	3u	1r	2r	T1	T2

	Expected coverage	Achievement (u)	Merit (r)	Excellence (t)
THREE (a)	$\frac{dy}{dx} = \frac{(x^2 + 1)(-\operatorname{cosec}^2 x) - (\cot x)(2x)}{(x^2 + 1)^2}$	Correct derivative.		
(b)	$\frac{dy}{dx} = \frac{2}{\sqrt{x}} - 1$ At stationary point, derivative = 0. $\frac{2}{\sqrt{x}} = 1$ $x = 4$ Coordinates are (4,6).	Correct solution with correct derivative.		
(c)	$\frac{dy}{dx} = \frac{(x^2 + 4) - x(2x)}{(x^2 + 4)^2}$ $= \frac{4 - x^2}{(x^2 + 4)^2}$ Increasing when $\frac{dy}{dx} > 0$ $\frac{4 - x^2}{(x^2 + 4)^2} > 0$ $4 - x^2 > 0$ $-2 < x < 2$	Correct $\frac{dy}{dx}$	Correct $\frac{dy}{dx}$ and identifies -2 and 2 as the boundaries of the interval required.	T1: Correct solution with correct derivative.
(d)	$\frac{dy}{dx} = \frac{(4x - k)4 - (4x + k)4}{(4x - k)^2}$ $= \frac{16x - 4k - 16x - 4k}{(4x - k)^2}$ $= \frac{-8k}{(4x - k)^2}$ When $x = 3$, $\frac{dy}{dx} = \frac{-8}{27}$ $\frac{-8k}{(12 - k)^2} = \frac{-8}{27}$ $\frac{k}{(12 - k)^2} = \frac{1}{27}$ $27k = 144 - 24k + k^2$ $k^2 - 51k + 144 = 0$ $k = 48$ or $k = 3$	Correct derivative.	Correct solution with correct derivative.	

<p>(e)</p> $\cos \theta = \frac{h}{S}$ $S^2 = h^2 + r^2$ $S = \sqrt{h^2 + r^2}$ <p>k and r are constant</p> $I = \frac{k \cos \theta}{S^2}$ $I = \frac{k \frac{h}{S}}{S^2}$ $= \frac{kh}{S^3}$ $I = \frac{kh}{(h^2 + r^2)^{\frac{3}{2}}}$ $\frac{dI}{dh} = \frac{(h^2 + r^2)^{\frac{3}{2}} k - kh \left(\frac{3}{2}\right) (h^2 + r^2)^{\frac{1}{2}} (2h)}{(h^2 + r^2)^3}$ $\frac{dI}{dh} = \frac{k(h^2 + r^2)^{\frac{3}{2}} - 3kh^2 (h^2 + r^2)^{\frac{1}{2}}}{(h^2 + r^2)^3}$ $\frac{dI}{dh} = \frac{k(h^2 + r^2)^{\frac{1}{2}} (h^2 + r^2 - 3h^2)}{(h^2 + r^2)^3}$ $\frac{dI}{dh} = \frac{k(r^2 - 2h^2)}{(h^2 + r^2)^{\frac{5}{2}}}$ $\frac{dI}{dh} = 0 \Rightarrow k(r^2 - 2h^2) = 0$ $2h^2 = r^2$ $h^2 = \frac{r^2}{2}$ $h = \frac{r}{\sqrt{2}}$		<p>Correct expression for $\frac{dI}{dh}$</p>	<p>T2: Correct proof with correct derivative</p>
---	--	--	--

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	ONE partial solution.	1u	2u	3u	1r	2r	T1	T2

Cut Scores

Not Achieved	Achievement	Achievement with Merit	Achievement with Excellence
0 – 6	7 – 12	13 – 18	19 – 24